




Evidence-Based Strategies for Supporting Older Adults with Depression & Anxiety

1

Today's Panelists



Colleen Toebe, MSN, CWCN, RAC-MT, RAC-MT, DNS MT
Vice President of Clinical Services
Pathway Health



Tiffany Robinson, BSW
Training Specialist
ComForCare Franchise Systems, LLC



2

Course Objectives

Upon completion of each session, participants should be able to:

- Identify common signs and risk factors for depression and anxiety in older adults.
- Understand appropriate screening and referral pathways.
- Compare pharmacologic and nonpharmacologic intervention strategies.
- Describe how engagement-focused programs support emotional well-being and cognitive health.



3





Common Signs and Symptoms

4

Depression in older adults

- Difficult to differentiate
- Irritable, may withdraw
- Can overlap with grief, dementia, chronic illness, pain, medications or sleep disruption.
- Loss of function, chronic, illness, pain and medications
- Change in the person's baseline





5

Mood changes

- Feeling sad, empty, hopeless or emotionally flat
- Increased tearfulness, pessimism, guilt, or feeling of worthlessness
- Irritability, frustration, anger, anxiety or restlessness



6

Loss of interest or social withdrawal

- No longer interested in activities, hobbies, meals, faith routines, or visitors
- Spending more time alone or wanting to stay in the room/home.
- Reduced conversation, eye contact, participation or initiation



7

Sleep pattern changes



- Trouble falling asleep, staying asleep, or early-morning waking
- Sleeping much more than usual or staying in bed during the day
- Fatigue after sleep, day-night reversal or increased napping

8

Appetite and weight changes

- Decreased appetite, skipped meals or food does not taste good.
- Unintentional weight loss- or overeating and weight gain
- Reduced fluid intake, poor motivation to eat or declining meal engagement



9

Low energy or functional decline

Fatigue, slowed movement, low motivation or "I just can't wait"

Decline in ADLs, hygiene, grooming, mobility or participation in therapy

Less initiation, waiting for others to prompt meals, toileting or activity

10

Cognitive and decision-making changes

- Trouble concentrating, remembering details, or making decisions
- Appearing confused, overwhelmed, indecisive or less engaged in choices
- Repeated negative statements such as "nothing matters" or "I'm a burden"
- Just want to die

11

Physical complaints or behavior changes

- Persistent aches, pain, headaches, stomach complaints or general discomfort
- Agitation, pacing, restlessness, repeated call light use or increased reassurance seeking
- Neglect of appearance, missed medications or decreased self-care



12

Safety signs



- Statement about suicide, self-harm, wanting to die or having no reason to live
- Giving away belongings, refusing essential care, or saying goodbye in an unusual way
- Sudden calm after distress, access to lethal means, or escalating hopelessness

13



Pharmacological and Non-Pharmacological Strategies


14

Medications


- Corticosteroids
- Benzodiazepines
- Sedative-hypnotics
- Hormonal therapy
- Digoxin
- Clonidine
- Methylidopa
- Reserpine
- Propranolol



15



16



17

- ### Caregiver stress
- Share the caregiving responsibilities
 - Use respite care
 - Maintain social connections
 - Protect sleep
 - See professional support
- ComForCare HOME CARE PATHWAY HEALTH

18

Sleep wake cycle

Corticosteroids

Stimulants

Decongestants

SSRI or SNRI

Beta Blockers

Thyroid

Bronchodilators

Cholinesterase Inhibitors

Benzodiazepines

ComForCare

PATHWAY HEALTH

19

Pharmacological interventions

Medication Class	Older Adult Benefit	Risks
Sertraline (Zoloft) – SSRI	Frequently used first-line; relatively few drug interactions; helpful for depression and anxiety	Nausea/diarrhea, hyponatremia, increased bleeding risk, falls, sexual dysfunction
Escitalopram (Lexapro) – SSRI	Often well tolerated; few interactions; useful when depression and anxiety occur together	Hyponatremia, falls, bleeding, GI effects
Citalopram (Celexa) – SSRI	Effective and generally well tolerated	Doses above 20 mg/day are generally avoided in adults over 60
Duloxetine (Cymbalta) – SNRI	May be particularly useful when depression occurs with neuropathic or chronic pain	Nausea, dizziness, blood-pressure effects, falls; renal/hepatic function must be considered
Venlafaxine (Effexor) – SNRI	Option when an SSRI is ineffective; treats depression and anxiety	May increase blood pressure; withdrawal symptoms if stopped abruptly; hyponatremia and falls
Mirtazapine (Remeron)	Can be helpful when depression occurs with poor appetite, weight loss, or insomnia	Sedation, increased appetite and weight gain, possible dizziness/falls
Bupropion (Wellbutrin)	Often less sedation, weight gain, and sexual dysfunction; may help low energy/apathy	May worsen insomnia or anxiety; can reduce appetite; avoid or use cautiously with seizure disorders

ComForCare

PATHWAY HEALTH

20

PATHWAY HEALTH

ComForCare

Anxiety in Older Adults

21

Anxiety in the older adult

- Often unrecognized
- 11.2 % of adults 65+ reported anxiety symptoms
- May appear as worry, fear, irritability or physical complaints
- Can overlap with depression
- Not normal aging
- Sleep disturbance

22

Common signs and symptoms

- Excessive worry and fear
- Restlessness, irritability, feeling on edge
- Poor sleep, fatigue
- Muscle tension, headaches, stomach upset
- Avoidance

23

Non-pharmacological interventions

- Cognitive behavioral therapy
- Relaxation
- Movement
- Connections
- Routine in daily activities

24

Medications

Medication	Examples	Related side effects
Decongestants – OTC	Pseudoephedrine (Sudafed), phenylephrine	Nervousness, restlessness, rapid heartbeat, insomnia
Caffeine-containing products – OTC	Some headache remedies, "alertness" products, energy products	Anxiety, tremor, irritability, insomnia, palpitations
Bronchodilators	Albuterol, ephedrine-containing products	Tremor, nervousness, rapid heartbeat, difficulty sleeping
Corticosteroids	Prednisone, methylprednisolone	Restlessness, mood changes, insomnia, agitation or anxiety
Thyroid replacement	Levothyroxine (Synthroid)	If the dose is too high: nervousness, anxiety, tremor, rapid heartbeat and insomnia
Some antidepressants	SSRIs/SNRIs, especially early in treatment	Temporary nervousness, activation, restlessness or insomnia in some people
Dopaminergic medications	Levodopa/carbidopa and some Parkinson's medications	Rebound anxiety, agitation, tremor and insomnia if reduced too quickly

New anxiety

- Did the anxiety begin after a medication was started
- Did the anxiety begin after a medication was increased
- Does the individual take cold, allergy or pain medication
- Has caffeine intake increased
- Did the person recently stop or reduce a sedative or anxiety

Mindfulness relaxation



Steps to mindful relaxation

- 5-minutes seated exercise
- Simple script
- Take slow deep breaths
- Short predictable and comfortable
- 3-5 minutes

Notice-Breath-Return




28




Screening in older adult for depression and anxiety

29

Assessment

- Ask about the onset, triggers, duration, severity and functional impact
- Screen for depression, grief, trauma, substance use and suicide risk
- Review medications, caffeine, alcohol, pain and sleep
- Consider cognition, delirium, sensory loss, falls and other safety concerns




30

Screening tools

Patient Name: _____ Date: _____

Instructions: Choose the best answer for how you feel over the past week. Note when asking the patient to complete the form, provide the self-rated form (included on the following page).

No. Question	Response	Score
1. Are you hardly satisfied with your life?	Yes/No	100/0
2. Have you stopped most of your activities and interests?	Yes/No	100/0
3. Do you feel that your life is empty?	Yes/No	100/0
4. Do you often get tired?	Yes/No	100/0
5. Do you go to your work or school often?	Yes/No	100/0
6. Do you feel that everything that is going to happen to you?	Yes/No	100/0
7. Do you feel things more of the time?	Yes/No	100/0
8. Do you often feel sad?	Yes/No	100/0
9. Do you prefer to stay at home rather than going out and doing other things?	Yes/No	100/0
10. Do you feel you have more problems with memory than usual?	Yes/No	100/0
11. Do you think it is worthwhile to go about?	Yes/No	100/0
12. Do you feel you are not the way you are now?	Yes/No	100/0
13. Do you feel that your situation is hopeless?	Yes/No	100/0
14. Do you feel that your situation is hopeless?	Yes/No	100/0
15. Do you feel that your situation is hopeless?	Yes/No	100/0
TOTAL:		

(David & Younger, 1980)

Scoring:
A score of 100 indicates depression is not evident; scores from 90 to 99 indicate a score of 100 is likely. A score greater than 1 suggests depression.

Notes:
• David & Younger (1980) Depression Scale (DS) is a 15-item measure of depression of a patient.
• Younger & David (1980) Depression Scale (DS) is a 15-item measure of depression of a patient.
• Younger & David (1980) Depression Scale (DS) is a 15-item measure of depression of a patient.
• Younger & David (1980) Depression Scale (DS) is a 15-item measure of depression of a patient.

Major Depressive Disorder Myths vs. Facts

Myth	Fact
Myth: Depression is not a medical condition.	Fact: Depression is a medical condition that can be treated with medication and therapy.
Myth: Depression is not a mental illness.	Fact: Depression is a mental illness that can be treated with medication and therapy.
Myth: Depression is not a chronic condition.	Fact: Depression is a chronic condition that can be treated with medication and therapy.
Myth: Depression is not a family illness.	Fact: Depression is a family illness that can be treated with medication and therapy.
Myth: Depression is not a life-threatening condition.	Fact: Depression is a life-threatening condition that can be treated with medication and therapy.

For Help and More Information:
For more information about mental health and wellness, visit www.comforcare.org or call 1-800-458-4584.

ComForCare HOME CARE **PATHWAY HEALTH**

31

Anxiety

CENTRAL ANXIETY DISORDER SCALE-7
RELATIVE RISK OF ANXIETY DISORDER SCALE-7

GAD-7

Over the last 7 weeks, how often have you been bothered by the following problems? (Use "0" to indicate your answer)


	Not at all	Several days	More than half the days	Nearly every day
1. Feeling nervous, anxious or on edge	0	1	2	3
2. Not being able to stop or control worrying	0	1	2	3
3. Worrying too much about different things	0	1	2	3
4. Trouble relaxing	0	1	2	3
5. Being so restless that it is hard to sit still	0	1	2	3
6. Becoming easily annoyed or irritable	0	1	2	3
7. Feeling afraid as if something awful might happen	0	1	2	3

(For office coding: Total Score 7 ____ * ____ * ____ * ____)

Developed by Drs. Robert L. Spitzer, Janet B.W. Williams, Kate Kessler and colleagues, with an educational grant from Pfizer Inc. No permission required to reproduce, translate, display or distribute.

ComForCare HOME CARE **PATHWAY HEALTH**

32



33



PATHWAY
HEALTH

High Education | Knowledge

ComForCare
HOME CARE

Training Strategies to Support Emotional & Cognitive Well-Being

34

Best practice for engagement-focused training

Training Care Teams to Put Meaningful Engagement Into Practice

- Embed **person-centered engagement principles** into training to help care teams understand individual interests, preferences, strengths, routines, and emotional needs.
- **Connect the "what" and the "why"** so teams understand how meaningful engagement can support emotional well-being, cognitive health, social connection, and quality of life.
- Use **real-world scenarios, role-playing, and case-based learning** to practice engaging older adults experiencing withdrawal, loss of interest, anxiety, or changes in participation.
- **Reinforce skills through coaching, mentoring, and feedback** to help teams personalize engagement and adapt approaches as needs and responses change.
- **Strengthen observation and communication skills** so care teams recognize meaningful changes and know when and how to share concerns with the appropriate care partners.



ComForCare
HOME CARE



PATHWAY
HEALTH

High Education | Knowledge

35

Creating consistency across care transitions

Supporting Emotional Well-Being Across Settings

- **Integrate depression and anxiety awareness into care-team training**, including changes in mood, behavior, participation, and social connection.
- **Build competency around transitions of care**, helping teams recognize how changes in health, function, environment, and support systems may affect emotional well-being.
- **Include person-centered engagement strategies** that can support connection, routine, purpose, and cognitive stimulation during and after transitions.
- **Establish clear communication, referral, and escalation pathways** so team members understand their role when concerns are identified.
- **Reinforce continuity across settings** by communicating successful approaches, preferences, observations, and changes from baseline.



ComForCare
HOME CARE



PATHWAY
HEALTH

High Education | Knowledge

36

12

Training in action: case scenario

Meet Mrs. Allen, age 81

- Hospitalized following a **hip fracture** requiring emergency **internal fixation (pinning)**
- Completed **21 days of skilled rehabilitation** with PT, OT, nursing, and mobility support
- History of **depression**; during rehab, staff noted periods of withdrawal and reduced motivation
- During rehabilitation, music, conversation, and staff interaction provided meaningful opportunities for engagement.
- Now transitioning home with **skilled home health and non-medical personal care support**
- Lives alone; daughter provides regular support



ComForCare HOME CARE PATHWAY HEALTH

37

Training in action: Applying engagement-focused support

1 Prepare the Team	2 Practice the Response	3 Reinforce Communication
<ul style="list-style-type: none"> Baseline & depression history Preferences, routines & what works Transition-home considerations Role & scope 	<ul style="list-style-type: none"> Withdrawal or increased anxiety Fear of falling / reduced participation Encourage without forcing Routine, choice & connection 	<ul style="list-style-type: none"> Observe & document changes Know who to notify and when Follow referral/escalation pathways Share what works across transition

TRAIN → PRACTICE → REINFORCE

Is the team prepared to recognize change, respond within role, and communicate concerns?

ComForCare HOME CARE PATHWAY HEALTH

38

Meaningful engagement across the continuum

Know what matters	Learn individual preferences, interests, routines, strengths, and what brings joy and comfort.
Share what works	Pass along successful strategies, approaches, and engagement activities that help.
Communicate meaningful changes	Observe and report changes in mood, behavior, participation, or social connection.
Engage family & care partners	Include family and key care partners in the plan, share updates, and collaborate on support.
Reassess as needs change	Check in regularly, adjust approaches, and continue to personalize support over time.



ComForCare HOME CARE PATHWAY HEALTH

39

Closing thought...

Effective training turns awareness into action—helping care teams recognize change, create meaningful connection, and support emotional well-being.



40

Helpful Resources

- [Depression and Older Adults | National Institute on Aging](#)
- [Clinical Practice Guidelines for Management of Depression in Elderly – PMC](#)
- [Services for Older Adults Living at Home | National Institute on Aging](#)
- [Pharmacotherapy of major depressive disorder in older adults: from an evidence-informed stepwise algorithm to precision medicine | Neuropsychopharmacology](#)
- [Drug Interactions: What You Should Know | FDA](#)



41

Disclaimer

This presentation provided is copyrighted information of Pathway Health and ComForCare Franchise Systems, LLC. Please note the presentation date on the title page in relation to the need to verify any new updates and resources that were listed in this presentation.

This presentation is intended to be informational.
The information does not constitute either legal or professional consultation.
This presentation is not to be sold or reused without written authorization.



42